

# Package ‘assessor’

April 3, 2024

**Title** Assessment Tools for Regression Models with Discrete and Semicontinuous Outcomes

**Version** 1.1.0

**Description** Provides assessment tools for regression models with discrete and semicontinuous outcomes proposed in Yang (2023) <[doi:10.48550/arXiv.2308.15596](https://doi.org/10.48550/arXiv.2308.15596)>. It calculates the double probability integral transform (DPIT) residuals, constructs QQ plots of residuals and the ordered curve for assessing mean structures.

**License** MIT + file LICENSE

**Encoding** UTF-8

**RoxygenNote** 7.3.1

**URL** <https://jhlee1408.github.io/assessor/>

**BugReports** <https://github.com/jhlee1408/assessor/issues>

**Imports** tweedie, MASS, VGAM, np

**Suggests** pscl, statmod, rmarkdown, knitr, AER, faraway, testthat (>= 3.0.0)

**Config/testthat/edition** 3

**Depends** R (>= 2.10)

**LazyData** true

**NeedsCompilation** no

**Author** Lu Yang [aut],  
Jeonghwan Lee [cre, aut]

**Maintainer** Jeonghwan Lee <[lee03938@umn.edu](mailto:lee03938@umn.edu)>

**Repository** CRAN

**Date/Publication** 2024-04-03 04:23:02 UTC

## R topics documented:

MEPS . . . . .	2
ord_curve . . . . .	3

qqresid . . . . .	5
resid_2pm . . . . .	6
resid_disc . . . . .	8
resid_quasi . . . . .	11
resid_semiconti . . . . .	13
resid_zeroinfl . . . . .	15

<b>Index</b>	<b>17</b>
--------------	-----------

---

MEPS	<i>Healthcare expenditure data</i>
------	------------------------------------

---

### Description

Healthcare expenditure data set.

### Usage

MEPS

### Format

A data frame with 29784 rows and 29 variables:

EXP the aggregate annual office based expenditure per participants, semicontinuous outcomes

AGE Age

GENDER 1 if female

ASIAN 1 if Asian

BLACK 1 if Black

NORTHEAST 1 if Northeast

MIDWEST 1 if Midwest

SOUTH 1 if South

USC 1 if have usual source of care

COLLEGE 1 if college or higher degrees

HIGHSCH 1 if high school degree

MARRIED 1 if married

WIDIVSEP 1 if widowed or divorced or separated

FAMSIZE Family Size

HINCOME 1 if high income

MINCOME 1 if middle income

LINCOME 1 if low income

NPOOR 1 if near poor

POOR 1 if poor

FAIR 1 if fair  
 GOOD 1 if good  
 VGOOD 1 if very good  
 MNHPOOR 1 if poor or fair mental health  
 ANYLIMIT 1 if any functional or activity limitation  
 unemployed 1 if unemployed at the beginning of 2006  
 EDUCHEALTH 1 if education, health and social services  
 PUBADMIN 1 if public administration  
 insured 1 if is insured at the beginning of the year 2006  
 MANAGEDCARE 1 if enrolled in an HMO or a gatekeeper plan

### Source

<http://www.meps.ahrq.gov/mepsweb/>

---

ord\_curve

*Ordered curve for assessing mean structures*

---

### Description

Creates a plot to assess the mean structure of regression models. The plot compares the cumulative sum of the response variable and its hypothesized value. Deviation from the diagonal suggests the possibility that the mean structure of the model is incorrect.

### Usage

`ord_curve(model, thr)`

### Arguments

model            Regression model object (e.g., `lm`, `glm`, `glm.nb`, `polr`, `lm`)  
 thr                Threshold variable (e.g., predictor, fitted values, or variable to be included as a covariate)

### Details

The ordered curve plots

$$\hat{L}_1(t) = \frac{\sum_{i=1}^n [Y_i 1(Z_i \leq t)]}{\sum_{i=1}^n Y_i}$$

against

$$\hat{L}_2(t) = \frac{\sum_{i=1}^n [\hat{\lambda}_i 1(Z_i \leq t)]}{\sum_{i=1}^n \hat{\lambda}_i},$$

where  $\hat{\lambda}_i$  is the fitted mean, and  $Z_i$  is the threshold variable.

If the mean structure is correctly specified in the model,  $\hat{L}_1(t)$  and  $\hat{L}_2(t)$  should be close to each other.

If the curve is distant from the diagonal, it suggests incorrectness in the mean structure. Moreover, if the curve is above the diagonal, the summation of the response is larger than the fitted mean, which implies that the mean is underestimated, and vice versa.

The role of thr (threshold variable  $Z$ ) is to determine the rule for accumulating  $\hat{\lambda}_i$  and  $Y_i$ ,  $i = 1, \dots, n$  for the ordered curve. The candidate for thr could be any function of predictors such as a single predictor (e.g.,  $x_1$ ), a linear combination of predictor (e.g.,  $x_1+x_2$ ), or fitted values (e.g., `fitted(model)`). It can also be a variable being considered to be included in the mean function. If a variable leads to a large discrepancy between the ordered curve and the diagonal, including this variable in the mean function should be considered.

For more details, see the reference paper.

### Value

- x-axis:  $\hat{L}_1(t)$
- y-axis:  $\hat{L}_2(t)$

which are defined in Details.

### References

Yang, Lu. "Double Probability Integral Transform Residuals for Regression Models with Discrete Outcomes." arXiv preprint arXiv:2308.15596 (2023).

### Examples

```
## Binary example of ordered curve
n <- 500
set.seed(1234)
x1 <- rnorm(n, 1, 1)
x2 <- rbinom(n, 1, 0.7)
beta0 <- -5
beta1 <- 2
beta2 <- 1
beta3 <- 3
q1 <- 1 / (1 + exp(beta0 + beta1 * x1 + beta2 * x2 + beta3 * x1 * x2))
y1 <- rbinom(n, size = 1, prob = 1 - q1)

## True Model
model0 <- glm(y1 ~ x1 * x2, family = binomial(link = "logit"))
ord_curve(model0, thr = model0$fitted.values) # set the threshold as fitted values

## Missing a covariate
model1 <- glm(y1 ~ x1, family = binomial(link = "logit"))
ord_curve(model1, thr = x2) # set the threshold as a covariate
```

```
## Poisson example of ordered curve
n <- 500
set.seed(1234)
x1 <- rnorm(n)
x2 <- rnorm(n)
beta0 <- 0
beta1 <- 2
beta2 <- 1
lambda1 <- exp(beta0 + beta1 * x1 + beta2 * x2)

y <- rpois(n, lambda1)

## True Model
poismodel1 <- glm(y ~ x1 + x2, family = poisson(link = "log"))
ord_curve(poismodel1, thr = poismodel1$fitted.values)

## Missing a covariate
poismodel2 <- glm(y ~ x1, family = poisson(link = "log"))
ord_curve(poismodel2, thr = poismodel2$fitted.values)
ord_curve(poismodel2, thr = x2)
```

---

qqresid

*QQ-plots of DPIT residuals*


---

## Description

Makes a QQ-plot of the DPIT residuals calculated from [resid\\_disc\(\)](#), [resid\\_semiconti\(\)](#) or [resid\\_zeroinfl\(\)](#). The plot should be close to the diagonal if the model is correctly specified. Note that this function does not return residuals. To get both residuals and QQ-plot, use [resid\\_disc\(\)](#), [resid\\_semiconti\(\)](#) and [resid\\_zeroinfl\(\)](#).

## Usage

```
qqresid(model, scale="normal")
```

## Arguments

model	Fitted model object (e.g., <a href="#">glm()</a> , <a href="#">glm.nb()</a> , <a href="#">zeroinfl()</a> , and <a href="#">polr()</a> )
scale	You can choose the scale of the residuals between normal and uniform scales. The sample quantiles of the residuals are plotted against the theoretical quantiles of a standard normal distribution under the normal scale, and against the theoretical quantiles of a uniform (0,1) distribution under the uniform scale. The default scale is normal.

## Value

A QQ plot.

- x-axis: Theoretical quantiles
- y-axis: Sample quantiles generated by DPIT residuals

**See Also**

`resid_disc()`, `resid_semiconti()`, `resid_zeroinfl()`

**Examples**

```
n <- 100
b <- c(2, 1, -2)
x1 <- rnorm(n)
x2 <- rbinom(n, 1, 0.7)
y <- rpois(n, exp(b[1] + b[2] * x1 + b[3] * x2))

m1 <- glm(y ~ x1 + x2, family = poisson)
qqresid(m1, scale = "normal")
qqresid(m1, scale = "uniform")
```

---

resid\_2pm

*Residuals for regression models with two-part outcomes*


---

**Description**

Calculates DPIT proposed residuals for model for semi-continuous outcomes. `resid_2pm` can be used either with `model0` and `model1` or with `part0` and `part1` as arguments.

**Usage**

```
resid_2pm(model0, model1, y, part0, part1, plot=TRUE, scale = "normal")
```

**Arguments**

<code>model0</code>	Model object for 0 outcomes (e.g., logistic regression)
<code>model1</code>	Model object for the continuous part (gamma regression)
<code>y</code>	Semicontinuous outcome variables
<code>part0</code>	Alternative argument to <code>model0</code> . One can supply the sequence of probabilities $P(Y_i = 0)$ , $i = 1, \dots, n$ .
<code>part1</code>	Alternative argument to <code>model1</code> . One can fit a regression model on the positive data and supply their probability integral transform. Note that the length of <code>part1</code> is the number of positive values in <code>y</code> and can be shorter than <code>part0</code> .
<code>plot</code>	A logical value indicating whether or not to return QQ-plot
<code>scale</code>	You can choose the scale of the residuals among normal and uniform scales. The default scale is normal.

## Details

The DPIT residuals for regression models with semi-continuous outcomes are

$$\hat{r}_i = \frac{\hat{F}(Y_i|\mathbf{X}_i)}{n} \sum_{j=1}^n 1 \left( \hat{p}_0(\mathbf{X}_j) \leq \hat{F}(Y_i|\mathbf{X}_i) \right), i = 1, \dots, n,$$

where  $\hat{p}_0(\mathbf{X}_i)$  is the fitted probability of zero, and  $\hat{F}(\cdot|\mathbf{X}_i)$  is the fitted cumulative distribution function for the  $i$ th observation. Furthermore,

$$\hat{F}(y|\mathbf{x}) = \hat{p}_0(\mathbf{x}) + (1 - \hat{p}_0(\mathbf{x})) \hat{G}(y|\mathbf{x})$$

where  $\hat{G}$  is the fitted cumulative distribution for the positive data.

In two-part models, the probability of zero can be modeled using a logistic regression, `model0`, while the positive observations can be modeled using a gamma regression, `model1`. Users can choose to use different models and supply the resulting probability transforms. `part0` should be the sequence of fitted probabilities of zeros  $\hat{p}_0(\mathbf{X}_i)$ ,  $i = 1, \dots, n$ . `part1` should be the probability integral transform of the positive part  $\hat{G}(Y_i|\mathbf{X}_i)$ . Note that the length of `part1` is the number of positive values in `y` and can be shorter than `part0`.

## Value

Residuals. If `plot=TRUE`, also produces a QQ plot.

## See Also

[resid\\_semiconti\(\)](#)

## Examples

```
library(MASS)
n <- 500
beta10 <- 1
beta11 <- -2
beta12 <- -1
beta13 <- -1
beta14 <- -1
beta15 <- -2
x11 <- rnorm(n)
x12 <- rbinom(n, size = 1, prob = 0.4)

p1 <- 1 / (1 + exp(-(beta10 + x11 * beta11 + x12 * beta12)))
lambda1 <- exp(beta13 + beta14 * x11 + beta15 * x12)
y2 <- rgamma(n, scale = lambda1 / 2, shape = 2)
y <- rep(0, n)
u <- runif(n, 0, 1)
ind1 <- which(u >= p1)
y[ind1] <- y2[ind1]

# models as input
mgamma <- glm(y[ind1] ~ x11[ind1] + x12[ind1], family = Gamma(link = "log"))
```

```

m10 <- glm(y == 0 ~ x12 + x11, family = binomial(link = "logit"))
resid.model <- resid_2pm(model0 = m10, model1 = mgamma, y = y)

# PIT as input
cdfgamma <- pgamma(y[ind1],
  scale = mgamma$fitted.values * gamma.dispersion(mgamma),
  shape = 1 / gamma.dispersion(mgamma)
)
p1f <- m10$fitted.values
resid.pit <- resid_2pm(y = y, part0 = p1f, part1 = cdfgamma)

```

resid\_disc

*Residuals for regression models with discrete outcomes***Description**

Calculates the DPIT residuals for regression models with discrete outcomes. Specifically, the model assumption of GLMs with binary, ordinal, Poisson, and negative binomial outcomes can be assessed using `resid_disc()`.

**Usage**

```
resid_disc(model, plot=TRUE, scale="normal")
```

**Arguments**

model	Model object (e.g., <code>glm</code> , <code>glm.nb</code> , <code>polr</code> )
plot	A logical value indicating whether or not to return QQ-plot
scale	You can choose the scale of the residuals among <code>normal</code> and <code>uniform</code> scales. The sample quantiles of the residuals are plotted against the theoretical quantiles of a standard normal distribution under the normal scale, and against the theoretical quantiles of a uniform (0,1) distribution under the uniform scale. The default scale is <code>normal</code> .

**Details**

The DPIT residual for the  $i$ th observation is defined as follows:

$$\hat{r}(Y_i|X_i) = \hat{G}\left(\hat{F}(Y_i|\mathbf{X}_i)\right)$$

where

$$\hat{G}(s) = \frac{1}{n-1} \sum_{j=1, j \neq i}^n \hat{F}\left(\hat{F}^{(-1)}(\mathbf{X}_j) \mid \mathbf{X}_j\right)$$

and  $\hat{F}$  refers to the fitted cumulative distribution function. When `scale="uniform"`, DPIT residuals should closely follow a uniform distribution, otherwise it implies model deficiency. When `scale="normal"`, it applies the normal quantile transformation to the DPIT residuals

$$\Phi^{-1}[\hat{r}(Y_i|\mathbf{X}_i)], i = 1, \dots, n.$$

The null pattern is the standard normal distribution in this case.

Check reference for more details.

## Value

DPIT residuals. If `plot=TRUE`, also produces a QQ plot.

## References

Yang, Lu. "Double Probability Integral Transform Residuals for Regression Models with Discrete Outcomes." arXiv preprint arXiv:2308.15596 (2023).

## Examples

```
library(MASS)
n <- 500
set.seed(1234)
## Negative Binomial example
# Covariates
x1 <- rnorm(n)
x2 <- rbinom(n, 1, 0.7)
### Parameters
beta0 <- -2
beta1 <- 2
beta2 <- 1
size1 <- 2
lambda1 <- exp(beta0 + beta1 * x1 + beta2 * x2)
# generate outcomes
y <- rnbinom(n, mu = lambda1, size = size1)

# True model
model1 <- glm.nb(y ~ x1 + x2)
resid.nb1 <- resid_disc(model1, plot = TRUE, scale = "uniform")

# Overdispersion
model2 <- glm(y ~ x1 + x2, family = poisson(link = "log"))
resid.nb2 <- resid_disc(model2, plot = TRUE, scale = "normal")

## Binary example
n <- 500
set.seed(1234)
# Covariates
x1 <- rnorm(n, 1, 1)
x2 <- rbinom(n, 1, 0.7)
# Coefficients
beta0 <- -5
beta1 <- 2
beta2 <- 1
beta3 <- 3
q1 <- 1 / (1 + exp(beta0 + beta1 * x1 + beta2 * x2 + beta3 * x1 * x2))
y1 <- rbinom(n, size = 1, prob = 1 - q1)
```

```

# True model
model01 <- glm(y1 ~ x1 * x2, family = binomial(link = "logit"))
resid.bin1 <- resid_disc(model01, plot = TRUE)

# Missing covariates
model02 <- glm(y1 ~ x1, family = binomial(link = "logit"))
resid.bin2 <- resid_disc(model02, plot = TRUE)

## Poisson example
n <- 500
set.seed(1234)
# Covariates
x1 <- rnorm(n)
x2 <- rbinom(n, 1, 0.7)
# Coefficients
beta0 <- -2
beta1 <- 2
beta2 <- 1
lambda1 <- exp(beta0 + beta1 * x1 + beta2 * x2)
y <- rpois(n, lambda1)

# True model
poismodel1 <- glm(y ~ x1 + x2, family = poisson(link = "log"))
resid.poi1 <- resid_disc(poismodel1, plot = TRUE)

# Enlarge three outcomes
y <- rpois(n, lambda1) + c(rep(0, (n - 3)), c(10, 15, 20))
poismodel2 <- glm(y ~ x1 + x2, family = poisson(link = "log"))
resid.poi2 <- resid_disc(poismodel2, plot = TRUE)

## Ordinal example
n <- 500
set.seed(1234)
# Covariates
x1 <- rnorm(n, mean = 2)
# Coefficient
beta1 <- 3

# True model
p0 <- plogis(1, location = beta1 * x1)
p1 <- plogis(4, location = beta1 * x1) - p0
p2 <- 1 - p0 - p1
genemult <- function(p) {
  rmultinom(1, size = 1, prob = c(p[1], p[2], p[3]))
}
test <- apply(cbind(p0, p1, p2), 1, genemult)
y1 <- rep(0, n)
y1[which(test[1, ] == 1)] <- 0
y1[which(test[2, ] == 1)] <- 1
y1[which(test[3, ] == 1)] <- 2
multimodel <- polr(as.factor(y1) ~ x1, method = "logistic")
resid.ord1 <- resid_disc(multimodel, plot = TRUE)

```

```

## Non-Proportionality
n <- 500
set.seed(1234)
x1 <- rnorm(n, mean = 2)
beta1 <- 3
beta2 <- 1
p0 <- plogis(1, location = beta1 * x1)
p1 <- plogis(4, location = beta2 * x1) - p0
p2 <- 1 - p0 - p1
genemult <- function(p) {
  rmultinom(1, size = 1, prob = c(p[1], p[2], p[3]))
}
test <- apply(cbind(p0, p1, p2), 1, genemult)
y1 <- rep(0, n)
y1[which(test[1, ] == 1)] <- 0
y1[which(test[2, ] == 1)] <- 1
y1[which(test[3, ] == 1)] <- 2
multimodel <- polr(as.factor(y1) ~ x1, method = "logistic")
resid.ord2 <- resid_disc(multimodel, plot = TRUE)

```

resid\_quasi

*Quasi Empirical residuals functions***Description**

Draw the QQ-plot for regression models with discrete outcomes using the quasi-empirical residual distribution functions. Specifically, the model assumption of GLMs with binary, ordinal, Poisson, negative binomial, zero-inflated Poisson, and zero-inflated negative binomial outcomes can be applicable to `resid_quasi()`.

**Usage**

```
resid_quasi(model)
```

**Arguments**

`model`                    Model object (e.g., `glm`, `glm.nb`, `polr`, `zeroinfl`)

**Details**

The quasi-empirical residual distribution function is defined as follows:

$$\hat{U}(s; \beta) = \sum_{i=1}^n W_n(s; \mathbf{X}_i, \beta) 1[F(Y_i | X_i) < H(s; X_i)]$$

where

$$W_n(s; \mathbf{X}_i, \beta) = \frac{K[(H(s; \mathbf{X}_i) - s)/\epsilon_n]}{\sum_{j=1}^n K[(H(s; \mathbf{X}_j) - s)/\epsilon_n]}$$

and  $K$  is a bounded, symmetric, and Lipschitz continuous kernel.

**Value**

A QQ plot.

- x-axis: Theoretical quantiles
- y-axis: Sample quantiles

**References**

Lu Yang (2021). Assessment of Regression Models with Discrete Outcomes Using Quasi-Empirical Residual Distribution Functions, *Journal of Computational and Graphical Statistics*, 30(4), 1019-1035.

**Examples**

```
## Negative Binomial example
library(MASS)
# Covariates
n <- 500
x1 <- rnorm(n)
x2 <- rbinom(n, 1, 0.7)
### Parameters
beta0 <- -2
beta1 <- 2
beta2 <- 1
size1 <- 2
lambda1 <- exp(beta0 + beta1 * x1 + beta2 * x2)
# generate outcomes
y <- rnbinom(n, mu = lambda1, size = size1)

# True model
model1 <- glm.nb(y ~ x1 + x2)
resid.nb1 <- resid_quasi(model1)

# Overdispersion
model2 <- glm(y ~ x1 + x2, family = poisson(link = "log"))
resid.nb2 <- resid_quasi(model2)

## Zero inflated Poisson example
library(pscl)
n <- 500
set.seed(1234)
# Covariates
x1 <- rnorm(n)
x2 <- rbinom(n, 1, 0.7)
# Coefficients
beta0 <- -2
beta1 <- 2
beta2 <- 1
beta00 <- -2
beta10 <- 2

# Mean of Poisson part
```

```

lambda1 <- exp(beta0 + beta1 * x1 + beta2 * x2)
# Excess zero probability
p0 <- 1 / (1 + exp(-(beta00 + beta10 * x1)))
## simulate outcomes
y0 <- rbinom(n, size = 1, prob = 1 - p0)
y1 <- rpois(n, lambda1)
y <- ifelse(y0 == 0, 0, y1)
## True model
modelzero1 <- zeroinfl(y ~ x1 + x2 | x1, dist = "poisson", link = "logit")
resid.zero1 <- resid_quasi(modelzero1)

```

resid\_semiconti

*Residuals for regression models with semicontinuous outcomes***Description**

Calculates the DPIT residuals for regression models with semi-continuous outcomes. The semi-continuous regression model such as a Tweedie regression model from `tweedie` package or a Tobit regression model from `VGAM`, `AER` packages is used in this function.

**Usage**

```
resid_semiconti(model, plot=TRUE, scale = "normal")
```

**Arguments**

<code>model</code>	Model object (e.g., <code>tweedie</code> , <code>vglm</code> , and <code>tobit</code> )
<code>plot</code>	A logical value indicating whether or not to return QQ-plot
<code>scale</code>	You can choose the scale of the residuals between normal and uniform scales. The default scale is normal.

**Details**

The DPIT residual for the  $i$ th semicontinuous observation is defined as follows:

$$\hat{r}_i = \frac{\hat{F}(Y_i|X_i)}{n} \sum_{j=1}^n I\left(\hat{p}_0(X_j) \leq \hat{F}(Y_i|X_i)\right),$$

which has a null distribution of uniformity.  $\hat{F}$  refers to the fitted cumulative distribution function, and  $\hat{p}_0$  refers to the fitted probability of being zero.

**Value**

Residuals. If `plot=TRUE`, also produces a QQ plot.

**References**

Lu Yang (2024). Diagnostics for Regression Models with Semicontinuous Outcomes, *Biometrics*, <https://arxiv.org/abs/2401.06347>

**See Also**

[resid\\_2pm\(\)](#)

**Examples**

```
## Tweedie model
library(tweedie)
library(statmod)
n <- 500
x11 <- rnorm(n)
x12 <- rnorm(n)
beta0 <- 5
beta1 <- 1
beta2 <- 1
lambda1 <- exp(beta0 + beta1 * x11 + beta2 * x12)
y1 <- rtweedie(n, mu = lambda1, xi = 1.6, phi = 10)
# Choose parameter p
# True model
model1 <-
  glm(y1 ~ x11 + x12,
      family = tweedie(var.power = 1.6, link.power = 0)
  )
resid.tweedie <- resid_semiconti(model1)

## Tobit regression model
library(VGAM)
beta13 <- 1
beta14 <- -3
beta15 <- 3

set.seed(1234)
x11 <- runif(n)
x12 <- runif(n)
lambda1 <- beta13 + beta14 * x11 + beta15 * x12
sd0 <- 0.3
yun <- rnorm(n, mean = lambda1, sd = sd0)
y <- ifelse(yun >= 0, yun, 0)

# Using VGAM package
# True model
fit1 <- vglm(formula = y ~ x11 + x12, tobit(Upper = Inf, Lower = 0, lmu = "identitylink"))
# Missing covariate
fit1miss <- vglm(formula = y ~ x11, tobit(Upper = Inf, Lower = 0, lmu = "identitylink"))

resid.tobit1 <- resid_semiconti(fit1, plot = TRUE)
resid.tobit2 <- resid_semiconti(fit1miss, plot = TRUE)

# Using AER package
library(AER)
# True model
fit2 <- tobit(y ~ x11 + x12, left = 0, right = Inf, dist = "gaussian")
# Missing covariate
```

```
fit2miss <- tobit(y ~ x11, left = 0, right = Inf, dist = "gaussian")
resid.aer1 <- resid_semiconti(fit2, plot = TRUE)
resid.aer2 <- resid_semiconti(fit2miss, plot = TRUE)
```

---

resid\_zeroinfl      *Residuals for regression models with zero-inflated outcomes*

---

### Description

Calculates the DPIT residuals for a regression model with zero-inflated discrete outcome. A zero-inflated model from `pscl` is used in this function.

### Usage

```
resid_zeroinfl(model, plot=TRUE, scale='normal')
```

### Arguments

<code>model</code>	Model object, which is the output of <code>pscl::zeroinfl</code> .
<code>plot</code>	A logical value indicating whether or not to return QQ-plot.
<code>scale</code>	You can choose the scale of the residuals among <code>normal</code> and <code>uniform</code> scales. The default scale is <code>normal</code> .

### Value

DPIT residuals. If `plot=TRUE`, also produces a QQ plot.

### References

Yang, Lu. "Double Probability Integral Transform Residuals for Regression Models with Discrete Outcomes." arXiv preprint arXiv:2308.15596 (2023).

### Examples

```
## Zero-Inflated Poisson
library(pscl)
n <- 500
set.seed(1234)
# Covariates
x1 <- rnorm(n)
x2 <- rbinom(n, 1, 0.7)
# Coefficients
beta0 <- -2
beta1 <- 2
beta2 <- 1
beta00 <- -2
beta10 <- 2

# Mean of Poisson part
```

```
lambda1 <- exp(beta0 + beta1 * x1 + beta2 * x2)
# Excess zero probability
p0 <- 1 / (1 + exp(-(beta00 + beta10 * x1)))
## simulate outcomes
y0 <- rbinom(n, size = 1, prob = 1 - p0)
y1 <- rpois(n, lambda1)
y <- ifelse(y0 == 0, 0, y1)
## True model
modelzero1 <- zeroinfl(y ~ x1 + x2 | x1, dist = "poisson", link = "logit")
resid.zero1 <- resid_zeroinfl(modelzero1, plot = TRUE, scale = "uniform")

## Zero inflation
modelzero2 <- glm(y ~ x1 + x2, family = poisson(link = "log"))
resid.zero2 <- resid_disc(modelzero2, plot = TRUE, scale = "normal")
```

# Index

## \* datasets

MEPS, 2

MEPS, 2

ord\_curve, 3

qqresid, 5

resid\_2pm, 6

resid\_2pm(), 14

resid\_disc, 8

resid\_disc(), 5, 6

resid\_quasi, 11

resid\_semiconti, 13

resid\_semiconti(), 5–7

resid\_zeroinfl, 15

resid\_zeroinfl(), 5, 6